

ENTIRE SOLUTIONS OF DONALDSON'S EQUATION

WEIYONG HE

ABSTRACT. We shall construct infinitely many special entire solutions to Donaldson's equation. We shall also prove a Liouville type theorem for entire solutions of Donaldson's equation. We believe that all entire solutions of Donaldson's equation have the form of the examples constructed in the paper.

1. INTRODUCTION

Donaldson [3] introduced an interesting differential operator when he set up a geometric structure for the space of volume forms on compact Riemannian manifolds. The Dirichlet problems for Donaldson's operator are considered in [2, 5]. In this note we shall consider this operator on Euclidean spaces.

For $(t, x) \in \Omega \subset \mathbb{R} \times \mathbb{R}^n$ ($n \geq 1$), let $u(t, x)$ be a smooth function such that $\Delta u > 0, u_{tt} > 0$. We use $\nabla u, \Delta u$ to denote derivatives with respect to x and $u_t = \partial_t u, u_{tt} = \partial_t^2 u$ to denote derivatives with respect to t . Define a differential operator Q by

$$Q(D^2u) = u_{tt}\Delta u - |\nabla u_t|^2.$$

This operator is strictly elliptic when $u_{tt} > 0, \Delta u > 0$ and $Q(D^2u) > 0$. When $n = 1$, then

$$Q(D^2u) = u_{tt}u_{xx} - u_{xt}^2$$

is a real Monge-Ampère operator. When $n = 2$, Q can be viewed as a special case of the complex Monge-Ampère operator. In x direction, we identify $\mathbb{R}^2 = \mathbb{C}$ with a coordinate w . In t direction, we take a product by $\mathbb{R}$ with a coordinate s and let $z = t + \sqrt{-1}s$. We extend u on $\mathbb{R} \times \mathbb{R}^2$ to $\mathbb{R}^4 = \mathbb{C}^2$ by $u(z, w) = u(t, x)$. Then

$$Q(D^2u) = 4(u_{z\bar{z}}u_{w\bar{w}} - u_{z\bar{w}}u_{w\bar{z}})$$

is a complex Monge-Ampère operator.

In this paper we shall consider entire solutions $u : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ of

$$(1.1) \quad Q(D^2u) = 1.$$

One celebrated result, proved by Jörgens [6] in dimension 2 and by Calabi [1] and Pogorelov [8] in higher dimensions, is that the only convex solutions of the real Monge-Ampère equation

$$(1.2) \quad \det(f_{ij}) = 1$$

on the whole of $\mathbb{R}^n$ are the obvious ones: quadratic functions.

Theorem 1.1 (Jörgens; Calabi; Pogorelov). *Let f be a global convex viscosity solution of (1.2) in whole $\mathbb{R}^n$. Then f has to be a quadratic function.*

2010 *Mathematics Subject Classification.* Primary 35J60, 35J96.

Key words and phrases. Donaldson's equation, entire solution.

The author is supported in-part by a start-up grant of University of Oregon.

One can also ask similar questions for the complex Monge-Ampère equations for plurisubharmonic functions. Let $v : \mathbb{C}^n \rightarrow \mathbb{R}$ be a strictly plurisubharmonic function such that $(v_{i\bar{j}}) > 0$, which satisfies

$$(1.3) \quad \det(v_{i\bar{j}}) = 1.$$

The similar results as Theorem 1.1 for the complex Monge-Ampère equation (1.3) or Donaldson's equation (1.1) ($n > 1$) are not known. For the complex Monge-Ampère equation, LeBrun [7] investigated the Euclidean Taub-NUT metric constructed by Hawking [4] and proved that it is a Kähler Ricci-flat metric on $\mathbb{C}^2$ but non-flat metric. His example provides a nontrivial entire solution of the complex Monge-Ampère equation. We shall construct infinitely many solutions for Donaldson's equation (1.1), which are nontrivial solutions in the sense that u_{tt} is a constant, but $\Delta u, \nabla u_t$ are both not constant. However, when $n = 2$, the Kähler metrics corresponding to these examples are the Euclidean metric on $\mathbb{C}^2$. We shall prove a Liouville type theorem for Donaldson's equation (1.1), which says u_{tt} has to be a constant provided some restriction on u_{tt} . Our proof relies on a transformation introduced by Donaldson [3]. We then ask if all solutions of (1.1) satisfy that u_{tt} is a constant; this would characterize all entire solutions of (1.1) if confirmed.

Acknowledgement: I am grateful to Prof. Xiuxiong Chen and Prof. Jingyi Chen for constant support and encouragements. I benefit from conversations with Prof. Pengfei Guan about the complex Monge-Ampère equations and I would like to thank him. I would also like to thank the referee for valuable suggestions and comments, and for spotting a minor error in the computation of the examples in Section 2 in a previous version of the paper.

2. EXAMPLES OF ENTIRE SOLUTIONS

In this section we shall construct infinitely many nontrivial solutions of (1.1) and (1.3). First we consider (1.1). Let $u_{tt} = 2a$ for some $a > 0$ and $u(0, x) = g(x), u_t(0, x) = b(x)$. Then

$$(2.1) \quad u(t, x) = at^2 + tb(x) + g(x).$$

If u solves (1.1), then

$$2a(t\Delta b + \Delta g) - |\nabla b|^2 = 1.$$

It follows that

$$\Delta b = 0, \Delta g = \frac{1}{2a} (1 + |\nabla b|^2).$$

So we shall construct the examples as follows. Let $b = b(x_1, x_2, \dots, x_n)$ be a harmonic function in $\mathbb{R}^n$. Define

$$h(x) = (1 + |\nabla b|^2)/2a.$$

Consider the following equation for $g(x)$

$$(2.2) \quad \Delta g = h(x).$$

Note that we can write $g = b^2(x)/4a + f$ for some function f such that $\Delta f = 1$. We can summarize our results above as follows.

Theorem 2.1. *Let u be the form of (2.1) such that b is a harmonic function and g satisfies (2.2), then u is an entire solution of (1.1). Moreover, any entire solution of (1.1) with $u_{tt} = \text{const}$ has the form of (2.1).*

When $n = 2$, the examples also provide solutions of the complex Monge-Ampère equation (1.3). Actually let $u(z, w) : \mathbb{C}^2 \rightarrow \mathbb{R}$ be a solution of (1.3). If $u_{z\bar{z}} = a$ for some constant $a > 0$, it is not hard to derive that

$$(2.3) \quad u(z, w) = az\bar{z} + f(z, \bar{z}) + zb(w, \bar{w}) + \bar{z}\bar{b}(w, \bar{w}) + g(w, \bar{w})$$

such that

$$\frac{\partial^2 f}{\partial z \partial \bar{z}} = \frac{\partial^2 b}{\partial w \partial \bar{w}} = 0, \quad \frac{\partial^2 g}{\partial w \partial \bar{w}} = \frac{1}{a} \left(1 + \left| \frac{\partial b}{\partial \bar{w}} \right|^2 \right).$$

However these examples are all trivial solutions of the complex Monge-Ampère equation in the sense that the corresponding Kähler metrics are flat.

3. A LIOUVILLE TYPE THEOREM

In this section we shall prove a Liouville type result for solutions of (1.1). We shall describe a transformation introduced by Donaldson [3], which relates the solutions of (1.1) with harmonic functions. Using this transformation, Theorem 3.1 follows from the standard Liouville theorem for positive harmonic functions.

Theorem 3.1. *Let u be a solution of (1.1) with $u_{tt} > 0$. For any $x \in \mathbb{R}^n$, if $u_{tt}(t, x)dt^2$ defines a complete metric on $\mathbb{R} \times \{x\}$, then u_{tt} is a constant. In particular, it has the form of (2.1) such that b is a harmonic function and g satisfies (2.2).*

Proof. For any x fixed, let $z = u_t(t, x)$. Then $\Phi : (t, x) \rightarrow (z, x)$ gives a transformation since $u_{tt} > 0$ and the Jacobian of Φ is always positive. In particular, when $u_{tt}(x, t)dt^2$ is a complete metric on $\mathbb{R} \times \{x\}$, $\Phi : (t, x) \rightarrow (z, x)$ is a diffeomorphism from $\mathbb{R} \times \mathbb{R}^n$ to $\mathbb{R} \times \mathbb{R}^n$. For x fixed, there exists a unique $t = t(z, x)$ such that $z = u_t(t, x)$. Define a function $\theta(z, x) = t(z, x)$. We claim that θ is a harmonic function in $\mathbb{R} \times \mathbb{R}^n$. The identity $z = u_t(\theta, x)$ implies

$$\frac{\partial \theta}{\partial x_i} u_{tt} + u_{tx_i} = 0, \quad u_{tt} \frac{\partial \theta}{\partial z} = 1.$$

It then follows that

$$\begin{aligned} u_{tt} \frac{\partial^2 \theta}{\partial x_i^2} + 2u_{ttx_i} \frac{\partial \theta}{\partial x_i} + u_{ttt} \left(\frac{\partial \theta}{\partial x_i} \right)^2 + u_{tx_i x_i} &= 0, \\ u_{tt} \frac{\partial^2 \theta}{\partial z^2} + \frac{u_{ttt}}{u_{tt}} &= 0. \end{aligned}$$

We compute, if u solves (1.1),

$$\begin{aligned}
\Delta_{(z,x)}\theta &= \frac{\partial^2\theta}{\partial z^2} + \sum_i \frac{\partial^2\theta}{\partial x_i^2} \\
&= \frac{1}{u_{tt}} \left(-\frac{u_{ttt}}{u_{tt}^2} - \Delta u_t + 2 \sum_i \frac{u_{ttx_i}u_{tx_i}}{u_{tt}} - \sum_i \frac{u_{ttt}u_{tx_i}^2}{u_{tt}^2} \right) \\
&= \frac{1}{u_{tt}} \left(-\frac{u_{ttt}}{u_{tt}^2} (1 + \sum u_{tx_i}^2) - \Delta u_t + 2 \sum_i \frac{u_{ttx_i}u_{tx_i}}{u_{tt}} \right) \\
&= \frac{1}{u_{tt}} \left(-\frac{u_{ttt}\Delta u}{u_{tt}} - \Delta u_t + 2 \sum_i \frac{u_{ttx_i}u_{tx_i}}{u_{tt}} \right) \\
&= -\frac{1}{u_{tt}^2} \partial_t (\Delta u u_{tt} - |\nabla u_t|^2) \\
&= 0.
\end{aligned}$$

On the other hand, $\partial\theta/\partial z = 1/u_{tt} > 0$. Hence $\partial\theta/\partial z$ is a positive harmonic function on $\mathbb{R} \times \mathbb{R}^n$. It follows that $\partial\theta/\partial z$ is a constant, and so u_{tt} is a constant. $\square$

One could classify all solutions of (1.1) if one could prove that u_{tt} decays to zero not too fast when $|t| \rightarrow \infty$ such that $u_{tt}dt^2$ defines a complete metric on a line. This motivates the following,

Problem 3.2. *Do all solutions of (1.1) with $u_{tt} > 0$ satisfy $u_{tt} = \text{const}$?*

REFERENCES

1. E. Calabi; *Improper affine hyperspheres of convex type and a generalization of a theorem by K. Jörgens*. Michigan Math. J. **5** (1958), 105–126.
2. X. Chen, W. He; *The space of volume forms*, arXiv:0810.3880.
3. S. K. Donaldson, *Nahm's equation and free boundary problems*, arXiv:0709.0184.
4. S. Hawking, *Gravitational instantons*, Phys. Lett. **60A** (1977), 8183.
5. W. He; *The Donaldson equation*, arXiv:0810.4123.
6. K. Jörgens; *Über die Lösungen der Differentialgleichung $rt - s^2 = 1$* . (German) Math. Ann. **127** (1954), 130–134.
7. C. LeBrun; *Complete Ricci-flat Kähler metrics on C^n need not be flat*. Several complex variables and complex geometry, Part 2 (Santa Cruz, CA, 1989), 297–304, Proc. Sympos. Pure Math., **52**, Part 2, Amer. Math. Soc., Providence, RI, 1991.
8. A.V. Pogorelov; *The Minkowski multidimensional problem*. Translated from the Russian by Vladimir Oliker. Introduction by Louis Nirenberg. Scripta Series in Mathematics. V. H. Winston & Sons, Washington, D.C.; Halsted Press [John Wiley & Sons], New York-Toronto-London, 1978.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OREGON, 97403
E-mail address: `whe@uoregon.edu`